

# Learning material

Site: [BOKUlearn - Die Lernplattform der Universität für Bodenkultur Wien](#)  
Course: 857304 Remote sensing and image processing (in Eng.) (VU) - 2022S  
Book: Learning material

Printed by: Vuolo Francesco  
Date: Thursday, 10 November 2022, 12:39 PM

# Table of contents

## **1. Introduction**

## **2. Semi-empirical reflectance model for LAI estimation**

## **3. Case study**

3.1. Description of the case study

3.2. R scripts, data and examples

## **4. References**

# 1. Introduction

The use of satellite sensor data has become more common in precision agriculture technologies (Lee et al. 2010), and various operational services are being developed within the Copernicus initiative (NEREUS 2012). In this context, the data users (i.e., farmers, large and small scale agri-businesses) are mostly interested in monitoring the spatial distribution of some crop characteristics over the growing season and time-series of satellite acquisitions at high spatial resolution are a major source of information.

A key vegetation parameter attracting most interest is the Leaf Area Index (LAI), defined as the total one-sided area of green leaf area per unit ground surface area (Bréda 2003). LAI is used to derive agronomic indicators for various crop management purposes. For instance, LAI maps are used in agro-meteorological models to derive the crop water needs (an example of operative application is given in Irrisat (D'Urso et al. 2010)), to monitor the nitrogen status and to apply fertilizer with variable rates (e.g., FarmSat), as input in crop models to derive agronomic variables (Jégo, Pattey, and Liu 2012; Casa et al. 2012). On a larger scale, LAI and other biophysical variables are used for example for yield predictions at administrative level (Rembold et al. 2012; Doraiswamy et al. 2005; Ma et al. 2013). A general overview of remote sensing contributions to agriculture is given in (Atzberger 2012).

Two groups of techniques have been commonly applied for the estimation of the LAI from optical satellite sensor data using semi-empirical/statistical approaches (i.e., vegetation indices, VI) or physical based approaches of leaf-canopy radiative transfer model (RTM) inversion (Baret and Buis 2008; Dorigo et al. 2007). Most of the empirical or statistical equations, such as regressions between spectral reflectance, vegetation indices (VI) or shape indices (e.g., red edge) and field measurements (Atzberger et al. 2010; Cho, Skidmore, and Atzberger 2008; Darvishzadeh et al. 2009), employ data in two or more wavebands, usually red and near-infrared (Chen et al. 2002; Y. J. Wang et al. 2004). VIs are often the only option for the retrieval of LAI with limited spectral information.

A prerequisite for the quantitative analysis of time-series of satellite sensor data is to perform radiometric and atmospheric corrections (Liu, Pattey, and Jégo 2012), if possible using reliable instantaneous atmospheric measurements (such as aerosol optical thickness, water vapor content) and/or the spectral reflectance of known ground targets either derived from ground measurements (surface and/or atmospheric conditions) or from consolidated library data (Hadjimitsis, Clayton, and Retalis 2009; R. Richter 1998). Several approaches have been proposed for performing atmospheric corrections. An operative procedure is based on the use of look-up-tables (LUT) with pre-calculated atmospheric RTM simulations for different satellite sensor types (R. Richter 1998; Liang, Fang, and Chen 2001). However, model assumptions and simplifications often lead to inaccuracies in the estimated top-of-canopy (TOC) reflectance measurements (Laurent et al. 2011).

For the implementation of this case study, we will use a simple LAI retrieval approach based on a VI using the CLAIR model (Clevers 1989). This approach has been tested using canopy reflectance model data (Clevers 1989), field-based reflectance measurements (Clevers et al. 2000) and satellite data in a number of studies (D'Urso et al. 2010; Vuolo et al., n.d.).

## 2. Semi-empirical reflectance model for LAI estimation

The CLAIR model is based on an inverse exponential relationship between LAI and the WdVI, which was derived using field spectral radiometer data experimentally (Clevers 1988, 1989). This relationship is built upon a simplified reflectance model of the light extinction through the vegetation canopy. The WdVI is calculated from the reflectance in red ( $\rho_{red}$ ) and near-infrared ( $\rho_{nir}$ ) as follows:

$$WdVI = \frac{\rho_{nir} - \rho_{red}}{\rho_{nir} + \rho_{red}} \quad \text{eq. 1}$$

where  $\rho_{nir} / \rho_{red}$  represents the 'soil line slope', a linear relationship between red and near-infrared reflectance of bare soils (Baret, Jacquemoud, and Hanocq 1993). The soil line slope parameter accounts for the effects of the soil background on the calculation of the vegetation index and it has to be determined for each test site (Baret and Guyot 1991).

Based on the WdVI, LAI is determined according to the following equation (Clevers 1988):

$$LAI = -1/\alpha \ln(1 - WdVI/WdVI_{\infty}) \quad \text{eq. 2}$$

where  $\alpha$  is a combination of extinction and scattering coefficients and  $WdVI_{\infty}$  is the asymptotically limiting value of WdVI for  $LAI \rightarrow \infty$ .

The  $\alpha$  coefficient can be estimated empirically using a set of LAI values from field measurements and contemporaneous reflectance values from satellite sensor data (Clevers 1989).

Generally,  $WdVI_{\infty}$  is derived directly from the image data considering the maximum WdVI value for vegetated areas in correspondence of saturation (dense vegetation with an expected LAI value of approximately 6-8 for cropland). Pasqualotto et al (2019) derived this value automatically by considering the mean ( $WdVI_{mean}$ ) and standard deviation ( $WdVI_{SD}$ ) as follows:

$$WdVI_{\infty} = WdVI_{mean} + 6WdVI_{SD} \quad \text{eq. 3}$$

The  $\alpha$  coefficient can be calibrated using a regression analysis technique applied to observed and estimated LAI values (D'Urso et al. 2010). This latter parameter describes the canopy architecture and it is dependent on the crop type and the corresponding Leaf Angle Distribution (LAD) value.

Different values of the parameters in Equation (2) can be found in the literature. For instance, Clevers (Clevers et al. 2000) reported an  $\alpha$  value ranging between 0.25 and 0.53 and  $WdVI_{\infty}$  ranging between 69 and 58 for vegetative and generative growth stages of barley respectively.

### 3. Case study

For the demonstration of this case study, we prepared a selection of about 60 sample points of bare soils (obtained from visual inspection) and the reflectance of these points will be used for calculation of the soil line slope.

The soil line slope will then be used to calculate the WdVI.

The WdVI<sub>0</sub> value can be extracted from the image data using the histogram or the procedure of eq 3.

We will derive the  $\alpha$  coefficient using an unconstrained nonlinear optimization method, which minimizes the Root Mean Square Error (RMSE) (our cost function) between measurements and predictions of LAI values. Measurements of LAI will be provided (CSV file). An initial  $\alpha$  value of 0.35 will be used as first guess based on previous experimental data.

To quantify the model prediction accuracy, we will use the following statistical measures: RMSE, the coefficient of determination ( $R^2$ ) between in-situ measured and predicted LAI.

Considering the limited number of the field LAI data that are usually available from in-situ measurements, the validation of the model performance can be achieved by resampling the field dataset using a bootstrapping approaches (Steyerberg et al. 2001) in order to provide an unbiased estimation of the model accuracy (Richter et al. 2012).

This technique will be however not used in this exercise.

Instead, the in-situ LAI dataset will be divided in Training and Testing (50/50 or 80/20) data.

### 3.1. Description of the case study

The field campaign for the collection of LAI data took place from April to August 2016. Eight different crop types, distributed over 72 parcels, were monitored to represent the prevailing crop types in the study area (Vuolo et al. 2018).

The parcels include 33 ordinary fields and 39 one-hectare experimental plots. In-situ LAI measurements were collected with a Li-Cor LAI-2200 Plant Canopy Analyzer (Li-Cor Inc. 2011).

The LAI-2200's sensor operates a non-destructive method and is sensitive to all light blocking objects in its view. It estimates the LAI from the values of canopy transmittance by identifying the attenuation of the radiation as it passes through the canopy (Li-Cor Inc. 2011). Therefore, measurements were taken above- (A) and below-canopy (B).

LAI estimates represent the effective Plant Area Index (PAI<sub>e</sub>), because the optical sensor does not distinguish between photosynthetically active leaves and inactive parts of the plants such as senescent leaves or stems. Care was taken to measure LAI only on photosynthetically active vegetation. For example, measurements were interrupted on winter cereals as soon as the first signs of senescence started to appear.

The LAI-2200 was deployed in elementary sampling units (ESUs) using a radius of 5-10 m of a geo-referenced point (accuracy of  $\pm 3$ -5m). Each unit represented a homogenous area with a single crop type. The ESUs were randomly chosen from the study area, with the only restriction being the fields' accessibility for time restraints. For ordinary fields the centres of the ESUs were placed in a corner of a squared area of 60m by 60m within the field and measured from the field border. It was imperative that the field conditions were relatively homogenous in terms of crop development. The ESUs located in the experimental plots, part of a larger experimental setup, were located in the centre of each one-hectare plot.

Winter cereal, onion, and potato were assessed through three replications of one A and eight B measurements, randomly distributed in the ESU. Row crops like maize, carrot and sugar beet were estimated with four replications of one A and six B measurements, for a total of 24 single measurements to generate a single LAI value per ESU (Weng 2017).

The final dataset of in-situ collected LAI consists of about 100 measurements and the corresponding L2A reflectance values extracted from the satellite image data. This dataset is made available for the execution of the case study.

## 3.2. R scripts, data and examples

### Materials:

- R Markdown document (LAI\_modelling.Rmd) containing the R-based script to be used in the exercise;
- In-Situ LAI measurements (file: LAI\_dataset.csv, column: LAI\_insitu);
- LAI obtained with an Artificial Neural Network not requiring local calibration (file: LAI\_dataset.csv, column: LAI\_S2\_BOKU);
- corresponding Level 2A (atmospherically corrected) reflectance (file: LAI\_dataset.csv, column name and spectral region):

1. B02 (490nm)
2. B03(560nm)
3. B04(665nm)
4. B05(705nm)
5. B06(740nm)
6. B07(783nm)
7. B08 (842nm)
8. B08A(865nm)
9. B11(1610nm)
10. B12(2190nm))

## 4. References

- Atzberger, Clement. 2012. "Advances in Remote Sensing of Agriculture: Context Description, Existing Operational Monitoring Systems and Major Information Needs." *Remote Sensing*, no. this issue.
- Atzberger, Clement, Martine Guérif, Frédéric Baret, and Willy Werner. 2010. "Comparative Analysis of Three Chemometric Techniques for the Spectroradiometric Assessment of Canopy Chlorophyll Content in Winter Wheat." *Computers and Electronics in Agriculture* 73 (2): 165–73. <https://doi.org/10.1016/j.compag.2010.05.006>.
- Baret, Frédéric, and Samuel Buis. 2008. "Estimating Canopy Characteristics from Remote Sensing Observations: Review of Methods and Associated Problems." In *Advances in Land Remote Sensing*. New York, USA., 173–201. wos:000255865000007.
- Baret, Frédéric, and G. Guyot. 1991. "Potentials and Limits of Vegetation Indices for LAI and APAR Assessment." *Remote Sensing of Environment* 35 (2–3): 161–73. [https://doi.org/10.1016/0034-4257\(91\)90009-U](https://doi.org/10.1016/0034-4257(91)90009-U).
- Baret, Frédéric, S. Jacquemoud, and J. F. Hanocq. 1993. "The Soil Line Concept in Remote Sensing." *Remote Sensing Reviews* 7 (1): 65–82. <https://doi.org/10.1080/02757259309532166>.
- Bonfante, A, MS Alfieri, A. Basile, F. De Lorenzi, N. Fiorentino, and M. Menenti. 2012. "The Evaluation of the Climate Change Effects on Maize and Fennel Cultivation by Means of a Hydrological Physically Based Model: The Case Study of an Irrigated District." *EGU General ...* 14: 8608. <http://adsabs.harvard.edu/abs/2012EGUGA..14.7807B>.
- Bréda, Nathalie J. J. 2003. "Ground-Based Measurements of Leaf Area Index: A Review of Methods, Instruments and Current Controversies." *Journal of Experimental Botany* 54 (392): 2403–17. <https://doi.org/10.1093/jxb/erg263>.
- Breiman, Leo. 2001. "Random Forests." *Machine Learning* 45 (1): 5–32.
- Camps-Valls, Gustavo, J. Munoz-Mari, L. Gomez-Chova, K. Richter, and J. Calpe-Maravilla. 2009. "Biophysical Parameter Estimation With a Semisupervised Support Vector Machine." *IEEE Geoscience and Remote Sensing Letters* 6 (2): 248–52. <https://doi.org/10.1109/LGRS.2008.2009077>.
- Capodici, F., Guido D'Urso, and A. Maltese. 2012. "Investigating the Relationship between X-Band SAR Data from COSMOS-Skymed Satellite and NDVI for LAI Detection." *Remote Sensing*, no. This issue.
- Casa, R., H. Varella, Samuel Buis, M. Guérif, B. De Solan, and Frédéric Baret. 2012. "Forcing a Wheat Crop Model with LAI Data to Access Agronomic Variables: Evaluation of the Impact of Model and LAI Uncertainties and Comparison with an Empirical Approach." *European Journal of Agronomy* 37 (1): 1–10. <https://doi.org/10.1016/j.eja.2011.09.004>.
- Chen, Jing M., G. Pavlic, L. Brown, J. Cihlar, S. G. Leblanc, H. P. White, R. J. Hall, et al. 2002. "Derivation and Validation of Canada-Wide Coarse-Resolution Leaf Area Index Maps Using High-Resolution Satellite Imagery and Ground Measurements." *Remote Sensing of Environment* 80 (1): 165–84. <http://www.sciencedirect.com/science/article/pii/S0034425701003005>.
- Cho, M. a., a. K. Skidmore, and Clement Atzberger. 2008. "Towards Red-edge Positions Less Sensitive to Canopy Biophysical Parameters for Leaf Chlorophyll Estimation Using Properties Optique Spectrales Des Feuilles (PROSPECT) and Scattering by Arbitrarily Inclined Leaves (SAILH) Simulated Data." *International Journal of Remote Sensing* 29 (8): 2241–55. <https://doi.org/10.1080/01431160701395328>.
- Clevers, Jan G.P.W. 1988. "The Derivation of a Simplified Reflectance Model for the Estimation of Leaf Area Index." *Remote Sensing of Environment* 25 (1): 53–69. [https://doi.org/10.1016/0034-4257\(88\)90041-7](https://doi.org/10.1016/0034-4257(88)90041-7).
- ———. 1989. "Application of a Weighted Infrared-Red Vegetation Index for Estimating Leaf Area Index by Correcting for Soil Moisture." *Remote Sensing of Environment* 29 (1): 25–37. [https://doi.org/10.1016/0034-4257\(89\)90076-X](https://doi.org/10.1016/0034-4257(89)90076-X).
- Clevers, Jan G.P.W., O.W. Vonder, R.E.E. Jongschaap, J.F. Desprats, C. King, L. Prévot, and N. Bruquier. 2000. "A Semi-Empirical Approach for Estimating Plant Parameters within the RESEDA-Project." *Proceedings of International Archives of Photogrammetry and Remote Sensing XXXIII*: 272–79.
- Coppola, E., G.F. Capra, P. Odierna, S. Vacca, and a. Buondonno. 2010. "Lead Distribution as Related to Pedological Features of Soils in the Volturno River Low Basin (Campania, Italy)." *Geoderma* 159 (3–4): 342–49. <https://doi.org/10.1016/j.geoderma.2010.08.010>.
- D'Urso, Guido, K. Richter, A. Calera, M. A. Osann, R. Escadafal, J. Garatuza-Pajan, L. Hanich, A. Perdigão, J. B. Tapia, and Francesco Vuolo. 2010. "Earth Observation Products for Operational Irrigation Management in the Context of the PLEIADeS Project." *Agricultural Water Management* 98 (2): 271–82. <https://doi.org/10.1016/j.agwat.2010.08.020>.
- Darvishzadeh, Roshanak, Clement Atzberger, A. K. Skidmore, and A. A. Abkar. 2009. "Leaf Area Index Derivation from Hyperspectral Vegetation Indices and the Red Edge Position." *International Journal of Remote Sensing* 30 (23): 6199–6218. <https://doi.org/10.1080/01431160902842342>.
- Doraiswamy, Paul C., Thomas R. Sinclair, Steven Hollinger, Bakhty Akhmedov, Alan Stern, and John Prueger. 2005. "Application of MODIS Derived Parameters for Regional Crop Yield Assessment." *Remote Sensing of Environment* 97 (2): 192–202. <https://doi.org/10.1016/j.rse.2005.03.015>.
- Dorigo, W.A., R. Zurita-Milla, A.J.W. de Wit, J. Brazile, R. Singh, and M.E. Schaepman. 2007. "A Review on Reflective Remote Sensing and Data Assimilation Techniques for Enhanced Agroecosystem Modeling." *International Journal of Applied Earth Observation and Geoinformation* 9 (2): 165–93. <https://doi.org/10.1016/j.jag.2006.05.003>.
- Garrigues, Sébastien, N.V. V. Shabanov, K. Swanson, J.T. T. Morisette, Frédéric Baret, and R.B. B. Myneni. 2008. "Intercomparison and Sensitivity Analysis of Leaf Area Index Retrievals from LAI-2000, AccuPAR, and Digital Hemispherical Photography over Croplands." *Agricultural and Forest Meteorology* 148 (8–9): 1193–1209. <https://doi.org/10.1016/j.agrformet.2008.02.014>.
- Hadjimitsis, Diofantos G., C.R.I. Clayton, and A. Retalis. 2009. "The Use of Selected Pseudo-Invariant Targets for the Application of Atmospheric Correction in Multi-Temporal Studies Using Satellite Remotely Sensed Imagery." *International Journal of Applied Earth Observation and Geoinformation* 11 (3): 192–200. <https://doi.org/10.1016/j.jag.2009.01.005>.
- Jégo, Guillaume, Elizabeth Pattey, and Jianguo Liu. 2012. "Using Leaf Area Index, Retrieved from Optical Imagery, in the STICS Crop Model for Predicting Yield and Biomass of Field Crops." *Field Crops Research* 131 (May): 63–74. <https://doi.org/10.1016/j.fcr.2012.02.012>.
- Laurent, Valérie C.E., Wout Verhoef, Jan G.P.W. Clevers, and Michael E. Schaepman. 2011. "Estimating Forest Variables from Top-of-Atmosphere Radiance Satellite Measurements Using Coupled Radiative Transfer Models." *Remote Sensing of Environment* 115 (4): 1043–52. <http://dx.doi.org/10.1016/j.rse.2010.12.009>.
- Leblanc, S. G., and Jing M. Chen. 2001. "A Practical Scheme for Correcting Multiple Scattering Effects on Optical LAI Measurements." *Agricultural and Forest Meteorology* 110 (2): 125–39. wos:000173333700004.
- Lee, W.S., V. Alchanatis, C. Yang, M. Hirafuji, D. Moshou, and C. Li. 2010. "Sensing Technologies for Precision Specialty Crop Production." *Computers and Electronics in Agriculture* 74 (1): 2–33. <https://doi.org/10.1016/j.compag.2010.08.005>.
- LI-COR. 1992. "LAI-2000 Plant Canopy Analyzer Instruction Manual Lincoln." NE, USA, LICOR Inc.
- Liang, Shunlin, Hongliang Fang, and Mingzhen Chen. 2001. "Atmospheric Correction of Landsat ETM+ Land Surface Imagery. Part I. Methods." *IEEE Transactions of Geoscience and Remote Sensing* 39 (11): 2490–98. <https://doi.org/10.1109/36.964986>.
- Liaw, Andy, and Matthew Wiener. 2002. "Classification and Regression by Random Forest." *R News* 2 (December): 18–22. [ftp://131.252.97.79/Transfer/Treg/WFRE\\_Articles/Liaw\\_02\\_Classification and regression by randomForest.pdf](ftp://131.252.97.79/Transfer/Treg/WFRE_Articles/Liaw_02_Classification and regression by randomForest.pdf).
- Liu, Jianguo, Elizabeth Pattey, and Guillaume Jégo. 2012. "Assessment of Vegetation Indices for Regional Crop Green LAI Estimation from Landsat Images over Multiple Growing Seasons." *Remote Sensing of Environment* 123 (August): 347–58. <https://doi.org/10.1016/j.rse.2012.04.002>.
- Ma, Guannan, Jianxi Huang, Wenbin Wu, Jinlong Fan, Jinqu Zou, and Sijie Wu. 2013. "Assimilation of MODIS-LAI into the WOFOST Model for Forecasting Regional Winter Wheat Yield." *Mathematical and Computer Modelling* 58 (3–4): 634–43. <https://doi.org/10.1016/j.mcm.2011.10.038>.
- Maire, Gueric le, Claire Marsden, Yann Nouvellon, Clovis Grinand, Rodrigo Hakamada, José-Luiz Stape, and Jean-Paul Laclau. 2011. "MODIS NDVI Time-Series Allow the Monitoring of Eucalyptus Plantation Biomass." *Remote Sensing of Environment* 115 (10): 2613–25. <http://dx.doi.org/10.1016/j.rse.2011.05.017>.
- Michele, C. De, Francesco Vuolo, Guido D'Urso, L. Marotta, and K. Richter. 2009. "The Irrigation Advisory Program of Campania Region: From Research to Operational Support for the Water Directive in Agriculture." In . Stresa, Italy.
- NEREUS. 2012. "The Growing Use of GMES Across Europe's Regions." *Chemistry & ...* <http://esamultimedia.esa.int/multimedia/publications/NEREUS/>.
- Powell, Scott L., Warren B. Cohen, Sean P. Healey, Robert E. Kennedy, Gretchen G. Moisen, Kenneth B. Pierce, and Janet L. Ohmann. 2010. "Quantification of Live Aboveground Forest Biomass Dynamics with Landsat Time-Series and Field Inventory Data: A Comparison of Empirical Modeling Approaches." *Remote Sensing of Environment* 114 (5): 1053–68. <https://doi.org/10.1016/j.rse.2009.12.018>.
- Rembold, F., Clement Atzberger, O. Rojas, and I. Savin. 2012. "Using Low Resolution Satellite Imagery for Yield Prediction and Yield Anomaly Detection." *Remote Sensing*, no. this issue.
- Richter, K., P. Rischbeck, J. Eitzinger, W. Schneider, F. Suppan, and P. Weihs. 2008. "Plant Growth Monitoring and Potential Drought Risk Assessment by Means of Earth Observation Data." *International Journal of Remote Sensing* 29 (17–18): 4943–60. <https://doi.org/10.1080/01431160802036268>.
- Richter, Katja, Clement Atzberger, Tobias B. Hank, and Wolfram Mauser. 2012. "Derivation of Biophysical Variables from Earth Observation Data: Validation and Statistical Measures Observation Data: Validation and Statistical Measures." *Journal of Applied Remote Sensing* 6 (1): <https://doi.org/10.1117/1.JRS.6.063557>.
- Richter, R. 1998. "Correction of Satellite Imagery over Mountainous Terrain." *Applied Optics* 37 (18): 4004–15. wos:000074127900026.
- Sequino, V., R. Zucaro, D. D'Alterio, M. Valente, M. Frunzio, V. Iavarone, E. Belmare, and G. Serpico. 1999. "Stato Dell'irrigazione in Campania."
- Smola, A.J., and B. Schölkopf. 2004. "A Tutorial on Support Vector Regression." *Statistics and Computing* 14: 199–222. <http://www.springerlink.com/index/KM7KRM46802R2114.pdf>.

- Steyerberg, E W, F E Harrell, G J Borsboom, M J Eijkemans, Y Vergouwe, and J D Habbema. 2001. "Internal Validation of Predictive Models: Efficiency of Some Procedures for Logistic Regression Analysis." *Journal of Clinical Epidemiology* 54 (8): 774–81. <http://www.ncbi.nlm.nih.gov/pubmed/11470385>.
- Suykens, J A K, T Van Gestel, J De Brabanter, Bar De Moor, and J Vandewalle. 2002. *Least Squares Support Vector Machines*. Singapore: World Scientific Pub.Co.
- Vapnik, V. 1995. *The Nature of Statistical Learning Theory*. Springer-Verlag New York, Inc. <http://dl.acm.org/citation.cfm?id=211359>.
- Vapnik, Vladimir, Steven E Golowich, and Alex J Smola. 1997. "Support Vector Method for Function Approximation, Regression Estimation and Signal Processing." In *Advances in Neural Information Processing Systems 9 --- Proceedings of the 1996 Neural Information Processing Systems Conference (NIPS 1996)*, edited by Michael Mozer, Michael I Jordan, and Thomas Petsche, 281–87. Dever, CO, USA: MIT Press, Cambridge, MA, USA. <http://dblp.uni-trier.de/db/conf/nips/nipsN1996.html#VapnikGS96>.
- Verrelst, Jochem, Jordi Muñoz, Luis Alonso, Jesús Delegido, Juan Pablo Rivera, Gustavo Camps-Valls, and José Moreno. 2012. "Machine Learning Regression Algorithms for Biophysical Parameter Retrieval: Opportunities for Sentinel-2 and -3." *Remote Sensing of Environment* 118 (March): 127–39. <https://doi.org/10.1016/j.rse.2011.11.002>.
- Vuolo, Francesco, Clement Atzberger, K Richter, Guido D'Urso, and Jadunandan Dash. n.d. "Retrieval of Biophysical Vegetation Products from RapidEye Imagery ." In *ISPRS TC VII Symposium - 100 Years ISPRS*, edited by W Wagner and B Székely, 7A ed., XXXVIII:281–86. Vienna, Austria. [http://www.isprs.org/proceedings/XXXVIII/part7/a/pdf/281\\_XXXVIII-part7A.pdf](http://www.isprs.org/proceedings/XXXVIII/part7/a/pdf/281_XXXVIII-part7A.pdf).
- Wang, Fu-min, Jing-feng Huang, and Zhang-hua Lou. 2011. "A Comparison of Three Methods for Estimating Leaf Area Index of Paddy Rice from Optimal Hyperspectral Bands." *Precision Agriculture* 12 (3): 439–47. <https://doi.org/10.1007/s11119-010-9185-2>.
- Wang, Y J, C E Woodcock, W Buermann, P Stenberg, P Voipio, H Smolander, T Hame, et al. 2004. "Evaluation of the MODIS LAI Algorithm at a Coniferous Forest Site in Finland." *Remote Sensing of Environment* 91 (1): 114–27. wos:000222005900008.
- Welles, I J, and J M Norman. 1991. "Instrument for Indirect Measurement of Canopy Architecture." *Agronomy Journal* 83 (5): 818–25. biosis:PREV199192119517.
- Yang, Xiaohua, Jingfeng Huang, Yaoping Wu, Jianwen Wang, Pei Wang, Xiaoming Wang, and Alfredo R Huete. 2011. "Estimating Biophysical Parameters of Rice with Remote Sensing Data Using Support Vector Machines." *Science China. Life Sciences* 54 (3): 272–81. <https://doi.org/10.1007/s11427-011-4135-4>.